

THE MODULI SPACE OF REAL ABELIAN VARIETIES WITH LEVEL STRUCTURE

MARK GORESKEY¹ AND YUNG SHENG TAI²

ABSTRACT. The moduli space of principally polarized abelian varieties with real structure and with level $N = 4m$ structure (with $m \geq 1$) is shown to coincide with the set of real points of a quasi-projective algebraic variety defined over $\mathbb{Q}$, and to consist of finitely many copies of the quotient of the space $\mathbf{GL}(n, \mathbb{R})/\mathbf{O}(N)$ (of positive definite symmetric matrices) by the principal congruence subgroup of level N in $\mathbf{GL}(n, \mathbb{Z})$.

1. INTRODUCTION

Let $\mathfrak{h}_n = \mathbf{Sp}(2n, \mathbb{R})/\mathbf{U}(n)$ be the Siegel upperhalf space of rank n . The quotient space $\mathbf{Sp}(2n, \mathbb{Z}) \backslash \mathfrak{h}_n$ has three remarkable properties: (a) it is the moduli space of principally polarized abelian varieties, (b) it has the structure of a quasi-projective complex algebraic variety which is defined over the rational numbers $\mathbb{Q}$, and (c) it has a natural compactification (the Baily-Borel Satake compactification) which is defined over the rational numbers.

Now let $C_n = \mathbf{GL}(n, \mathbb{R})/\mathbf{O}(n)$ be the symmetric cone of positive definite symmetric matrices and let $Z = \mathbf{GL}(n, \mathbb{Z}) \backslash C_n$. One might ask whether similar statements hold for Z . Is Z in some sense a moduli space for principally polarized real abelian varieties? Does it admit the structure of a real algebraic variety, possibly defined over $\mathbb{Q}$? If so, does it admit a compactification which is also defined over the rational numbers? The answer to all these questions is “no”. In fact, Silhol [Si] constructs the moduli space of real principally polarized abelian varieties and he shows that it is a (topological) ramified covering of Z . Moreover Silhol constructs a compactification of this moduli space, analogous to the Baily-Borel compactification. However neither the moduli space nor this compactification has an algebraic structure.

In this paper we show that all three statements (a), (b), and (c) above may be recovered if we consider real abelian varieties with an appropriate level structure. To be precise, let $N = 4m$ for some $m \geq 1$, let $\Gamma_\ell(N)$ be the principal congruence subgroup of $\mathbf{GL}(n, \mathbb{Z})$ of level N , and let $Z(N) = \Gamma_\ell(N) \backslash C_n$. Then (a) the moduli space of real principally polarized abelian varieties with level N structure consists of finitely many copies of $Z(N)$. These various copies are indexed by a certain (nonabelian cohomology) set $H^1(\mathbb{C}/\mathbb{R}, \Gamma(N))$. Moreover, (b)

1. School of Mathematics, Institute for Advanced Study, Princeton N.J. Research partially supported by NSF grants # DMS 9626616 and DMS 9900324.

2. Dept. of Mathematics, Haverford College, Haverford PA..

this moduli space may be naturally identified with the real points $X_{\mathbb{R}}$ of a quasi-projective algebraic variety X which is defined over the rational numbers. See Theorems 4.4 and 6.6. Finally, (c) the variety $X_{\mathbb{C}}$ has a compactification which is defined over $\mathbb{Q}$.

In fact the variety $X_{\mathbb{C}}$ is just the moduli space $\Gamma(N) \backslash \mathfrak{h}_n$ of abelian varieties with principal polarization and level $N = 4m$ structure. (Here, $\Gamma(N)$ denotes the principal congruence subgroup of $\mathbf{Sp}(2n, \mathbb{Z})$ of level N .) The key observation is that there is an involution τ on $\mathbf{Sp}(2n, \mathbb{R})$ whose fixed point set is $\mathbf{GL}(n, \mathbb{R})$, which passes to an anti-holomorphic involution on $X_{\mathbb{C}}$.

Let $V = V_{\mathbb{C}}$ denote the Baily-Borel Satake compactification of $X_{\mathbb{C}}$. It is an algebraic variety defined over $\mathbb{Q}$. Complex conjugation $\tau : V_{\mathbb{C}} \rightarrow V_{\mathbb{C}}$ is an anti-holomorphic involution whose fixed point set is the set of real points $V_{\mathbb{R}}$. This set is compact and it contains $X_{\mathbb{R}}$ as an open set. One might ask whether $V_{\mathbb{R}}$ is a compactification of $X_{\mathbb{R}}$, that is, whether $V_{\mathbb{R}} = \overline{X}_{\mathbb{R}}$, where $\overline{X}_{\mathbb{R}}$ denotes the closure of $X_{\mathbb{R}}$ in $V_{\mathbb{C}}$. We have been able to prove this (§10) in the special case $N = 2^k$ (that is, for the principal congruence subgroup $\Gamma(2^k)$ of level 2^k) for $k \geq 2$, and we suspect it is otherwise false unless $n = 1$; cf §5. However we have been able to show (for general $\Gamma(4m)$) that the complement $V_{\mathbb{R}} - \overline{X}_{\mathbb{R}}$ has a high codimension. The variety $V_{\mathbb{C}}$ has a stratification

$$V_{\mathbb{C}} = V_{\mathbb{C}}^0 \cup V_{\mathbb{C}}^1 \cup \dots \cup V_{\mathbb{C}}^n$$

such that each $V_{\mathbb{C}}^r$ is a union of arithmetic quotients $\Gamma \backslash \mathfrak{h}_{n-r}$ of finitely many copies of the Siegel space of rank $n - r$. The largest stratum $V_{\mathbb{C}}^0$ is just $X_{\mathbb{C}}$. Denote by $V_{\mathbb{R}}^r$ the τ -fixed points in $V_{\mathbb{C}}^r$. In Proposition 9.2 we show that

$$V_{\mathbb{R}}^0 \cup V_{\mathbb{R}}^1 \subset \overline{X}_{\mathbb{R}} \subset V_{\mathbb{R}}.$$

Although the moduli space $X_{\mathbb{R}}$ consists of finitely many disjoint copies of the locally symmetric space $Z(N)$, the compactification $\overline{X}_{\mathbb{R}}$ is not a disjoint union: some of these copies of $Z(N)$ may become glued together along the boundary.

The authors are grateful to the Institute for Advanced Study in Princeton N.J. for its hospitality and support.

2. SYMPLECTIC GROUP

2.1. The symplectic group $G = \mathbf{Sp}(2n, \mathbb{R})$ may be realized as the group of $2n$ by $2n$ real matrices

$$\begin{aligned} \mathbf{Sp}(2n, \mathbb{R}) &= \left\{ \begin{pmatrix} A & B \\ C & D \end{pmatrix} \mid \begin{array}{l} {}^tAD - {}^tCB = I \\ {}^tAC, {}^tBD \text{ symmetric} \end{array} \right\} \\ &= \left\{ \begin{pmatrix} A & B \\ C & D \end{pmatrix} \mid \begin{array}{l} A {}^tD - B {}^tC = I \\ A {}^tB, C {}^tD \text{ symmetric} \end{array} \right\} \end{aligned}$$

If $J = \begin{pmatrix} 0 & I_n \\ -I_n & 0 \end{pmatrix}$ is the standard symplectic form on $\mathbb{R}^{2n}$ then $g \in \mathbf{Sp}(2n, \mathbb{R}) \iff {}^t g J g = J$.

The inverse of such a symplectic matrix is

$$\begin{pmatrix} A & B \\ C & D \end{pmatrix}^{-1} = \begin{pmatrix} {}^t D & -{}^t B \\ -{}^t C & {}^t A \end{pmatrix} \quad (2.1.1)$$

Identify $\mathbf{GL}(n, \mathbb{R}) \hookrightarrow \mathbf{Sp}(2n, \mathbb{R})$ with its image under the embedding

$$A \mapsto \begin{pmatrix} A & 0 \\ 0 & {}^t A^{-1} \end{pmatrix}$$

A Cartan involution θ of $\mathbf{Sp}(2n, \mathbb{R})$ is given by $\theta g = J g J^{-1}$, that is,

$$\theta \begin{pmatrix} A & B \\ C & D \end{pmatrix} = \begin{pmatrix} D & -C \\ -B & A \end{pmatrix}$$

Its fixed point set is the unitary group $K = \mathbf{U}(n)$ which is embedded in the symplectic group by

$$A + iB \mapsto \begin{pmatrix} A & B \\ -B & A \end{pmatrix}.$$

The symplectic group acts transitively on the Siegel upper halfspace

$$\mathfrak{h}_n = \left\{ Z = X + iY \in M_{n \times n}(\mathbb{C}) \mid {}^t Z = Z \text{ and } Y > 0 \right\}$$

by fractional linear transformations: if $g = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$ then

$$gZ = (AZ + B)(CZ + D)^{-1}.$$

Then $\text{Stab}_G(iI) = \mathbf{U}(n)$, and $\mathfrak{h}_n \cong G/K$.

2.2. Let $I_- = \begin{pmatrix} -I & 0 \\ 0 & I \end{pmatrix} \in \mathbf{GL}(2n, \mathbb{R})$. Consider the involution $\tau : \mathbf{Sp}(2n, \mathbb{R}) \rightarrow \mathbf{Sp}(2n, \mathbb{R})$ which is defined by $\tau(g) = I_- g I_-$, that is,

$$\tau \begin{pmatrix} A & B \\ C & D \end{pmatrix} = \begin{pmatrix} A & -B \\ -C & D \end{pmatrix} \quad (2.2.1)$$

The following properties of τ are easily verified by direct calculation:

1. τ is an involution of G , i.e. $\tau(gh^{-1}) = \tau(g)(\tau(h))^{-1}$.
2. $\tau(g) = g \iff g \in \mathbf{GL}(n, \mathbb{R})$
3. $\tau\theta = \theta\tau$ so $\tau(\mathbf{U}(n)) = \mathbf{U}(n)$
4. If $C = A + iB \in \mathbf{U}(n)$ then $\tau(C) = \bar{C} = A - iB$.

It follows that τ passes to an involution (which we denote by the same letter) $\tau : \mathfrak{h}_n \rightarrow \mathfrak{h}_n$ on the Siegel space such that

$$\tau(g \cdot Z) = \tau(g) \cdot \tau(Z)$$

for all $g \in \mathbf{Sp}(2n, \mathbb{R})$ and for all $Z \in \mathfrak{h}_n$. In fact $\tau : \mathfrak{h}_n \rightarrow \mathfrak{h}_n$ is the antiholomorphic involution given by

$$\tau(Z) = -\overline{Z}.$$

Its fixed point set is the orbit

$$iC_n = \mathbf{GL}(n, \mathbb{R}).iI \subset M_{n \times n}(\mathbb{C}) \quad (2.2.2)$$

of the general linear group. Here, $C_n \cong \mathbf{GL}(n, \mathbb{R})/\mathbf{O}(n)$ denotes the cone of positive definite symmetric real matrices.

2.3. Notation. For $\gamma \in G$ and $Z \in \mathfrak{h}_n$ we will usually write $\tilde{\gamma} = \tau(\gamma)$ and $\tilde{Z} = \tau(Z)$. For any $\gamma \in \mathbf{Sp}(2n, \mathbb{R})$, define the locus of γ -real points (or, more precisely, the set of (γ, τ) -real points),

$$\mathfrak{h}_n^\gamma = \{Z \in \mathfrak{h}_n \mid \gamma Z = \tau(Z)\} \quad (2.3.1)$$

If $\Gamma \subset \mathbf{Sp}(2n, \mathbb{R})$ is an arithmetic subgroup such that $\tau\Gamma = \Gamma$, define the set $\mathfrak{h}_n^\Gamma$ of Γ -real points to be the union $\bigcup_{\gamma \in \Gamma} \mathfrak{h}_n^\gamma$.

3. GALOIS COHOMOLOGY

3.1. The involution τ of (2.2.1) may be considered as an action of $\text{Gal}(\mathbb{C}/\mathbb{R}) = \{1, \tau\}$ on $\mathbf{Sp}(2n, \mathbb{R})$. Its restriction to the unitary group $K = \mathbf{U}(n) \subset \mathbf{Sp}(2n, \mathbb{R})$ coincides with complex conjugation. For any $\gamma \in \mathbf{Sp}(2n, \mathbb{R})$ let $f_\gamma : \text{Gal}(\mathbb{C}/\mathbb{R}) \rightarrow \mathbf{Sp}(2n, \mathbb{R})$ be the mapping $f_\gamma(1) = I$ and $f_\gamma(\tau) = \gamma$. Then f_γ is a 1-cocycle iff $\gamma\tilde{\gamma} = I$, and f_γ is a coboundary iff there exists $h \in \mathbf{Sp}(2n, \mathbb{R})$ so that $\gamma = \tilde{h}h^{-1}$. In this section we will prove the following:

3.2. Proposition. *With respect to the above action (2.2.1), the (nonabelian) Galois cohomology sets $H^1(\mathbb{C}/\mathbb{R}, \mathbf{U}(n))$ and $H^1(\mathbb{C}/\mathbb{R}, \mathbf{Sp}(2n, \mathbb{R}))$ are trivial.*

3.3. Lemma. *If $g \in \mathbf{U}(n)$ and $g\tilde{g} = I$ then there exists $\delta \in \mathbf{U}(n)$ such that $g = \tilde{\delta}\delta^{-1}$.*

3.4. Proof. For completeness we give a short proof of this well known result, which is the first part of Proposition 3.2. Set $g = \begin{pmatrix} A & B \\ -B & A \end{pmatrix}$. Calculating $g\tilde{g} = I$ gives $A^2 + B^2 = I$ and $AB = BA$. Comparing with (2.1.1) gives ${}^tA = A$ and ${}^tB = B$. Such commuting symmetric matrices may be simultaneously diagonalized by an orthogonal matrix $h \in \mathbf{O}(n)$. In other words, $h(A + iB)h^{-1} = \Lambda \in \mathbf{U}(n)$ is a diagonal matrix. Choose a square root, $\mu = \sqrt{\Lambda} \in \mathbf{U}(n)$ (by choosing a square root of each diagonal entry) and set $\delta = h^{-1}\mu h$. Since $\tilde{h} = h$ and $\tilde{\mu} = \mu^{-1}$, we find

$$\tilde{\delta}\delta^{-1} = h^{-1}\tau(\mu)\mu^{-1}h = h^{-1}\tilde{\mu}\tilde{\mu}h = h^{-1}\Lambda h = A + iB. \quad \square$$

3.5. Lemma. *Let $\gamma \in \mathbf{Sp}(2n, \mathbb{R})$ and suppose that f_γ is a 1-cocycle (i.e., $\gamma\tilde{\gamma} = I$). Then f_γ is a coboundary iff $\mathfrak{h}_n^\gamma \neq \emptyset$.*

3.6. Proof. Suppose f_γ is a coboundary, say $\gamma = \tilde{h}h^{-1}$. Then $\gamma hiY = \tau(hiY)$ for any $Y \in C_n$, so $\mathfrak{h}_n^\gamma \neq \phi$. On the other hand, suppose $Z \in \mathfrak{h}_n^\gamma$. Write $Z = h(iI)$ for some $h \in \mathbf{Sp}(2n, \mathbb{R})$. Then $\tilde{h}^{-1}\gamma h(iI) = \tilde{h}^{-1}\gamma Z = \tilde{h}^{-1}\tilde{Z} = \tau(iI) = iI$ hence $\alpha = \tilde{h}^{-1}\gamma h \in \mathbf{U}(n)$. Moreover, $\alpha\tilde{\alpha} = \tilde{h}^{-1}\gamma\tilde{\gamma}\tilde{h} = I$ so by Lemma 3.3 there exists $\delta \in \mathbf{U}(n)$ with $\alpha = \tilde{\delta}\delta^{-1}$. Therefore $\gamma = \tau(h\delta)(h\delta)^{-1}$. $\square$

3.7. Suppose that $\gamma = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \mathbf{Sp}(2n, \mathbb{R})$ and that $\gamma\tilde{\gamma} = I$. Then

$$A = {}^tD, \ C = {}^tC, \ B = {}^tB, \ A^2 - BC = I, \ CA = {}^tAC, \ \text{and} \ AB = B{}^tA. \quad (3.7.1)$$

3.8. Lemma. Suppose that $\gamma = \begin{pmatrix} 0 & -C^{-1} \\ C & 0 \end{pmatrix} \in \mathbf{Sp}(2n, \mathbb{R})$ and that C is diagonal. Then the cocycle f_γ is cohomologically trivial.

3.9. Proof. By Lemma 3.5 it suffices to show that there exists $Z \in \mathfrak{h}_n$ such that $\gamma \cdot Z = -C^{-1}Z^{-1}C^{-1} = -\bar{Z}$. If $C = \text{diag}(c_1, c_2, \dots, c_n)$ then a solution is given by $Z = iY$ where $Y = \text{diag}(y_1, y_2, \dots, y_n)$ and $y_i = |c_i|^{-1}$. $\square$

3.10. Lemma. Suppose $\gamma = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \mathbf{Sp}(2n, \mathbb{R})$. Suppose that $\gamma\tilde{\gamma} = I$ and that $C = 0$. Then the 1-cocycle f_γ is cohomologically trivial.

3.11. Proof. The cocycle f_γ is equivalent to $f_{\gamma'}$ where

$$\gamma' = \begin{pmatrix} A & 0 \\ 0 & {}^tA \end{pmatrix} = \tilde{h}\gamma h^{-1}$$

and

$$h = \begin{pmatrix} I & \frac{1}{2}A^{-1}B \\ 0 & I \end{pmatrix}$$

using (3.7.1). Then $A^2 = I$ so A is $\mathbf{GL}(n, \mathbb{R})$ -conjugate to a diagonal matrix Λ (with eigenvalues ± 1). Therefore $f_{\gamma'}$ is equivalent to the 1-cocycle defined by the element $\begin{pmatrix} \Lambda & 0 \\ 0 & \Lambda^{-1} \end{pmatrix} \in \mathbf{U}(n)$. It follows from Lemma 3.3 that $f_{\gamma'}$ is cohomologically trivial. $\square$

3.12. Proof of Proposition 3.2. Let $\gamma' = \begin{pmatrix} A' & B' \\ C' & D' \end{pmatrix}$ and suppose that $f_{\gamma'}$ is a cocycle. By (3.7.1), C' is symmetric so there exists $k \in \mathbf{O}(n)$ so that $C = kC'{}^tk = \begin{pmatrix} C_1 & 0 \\ 0 & 0 \end{pmatrix}$ where $C_1 \in M_{r \times r}(\mathbb{R})$ is diagonal and nonsingular. Let $h_1 = \begin{pmatrix} k & 0 \\ 0 & k \end{pmatrix}$. Then $f_{\gamma'}$ is cohomologous to f_γ where

$$\gamma = \tilde{h}_1\gamma'h_1^{-1} = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$$

and (3.7.1) holds. Write $A = \begin{pmatrix} A_1 & A_2 \\ A_3 & A_4 \end{pmatrix}$ where $A_1 \in M_{r \times r}(\mathbb{R})$ and $A_4 \in M_{(n-r) \times (n-r)}(\mathbb{R})$, and similarly for B and D . Then $CA = {}^tAC$ implies that $A_2 = A_3 = 0$.

Set $h_2 = \begin{pmatrix} I & x \\ 0 & I \end{pmatrix}$ where

$$x = \begin{pmatrix} C_1^{-1} {}^t A_1 & 0 \\ 0 & 0 \end{pmatrix}.$$

Then $A^2 - BC = I$ implies that f_γ is cohomologous to f_g where

$$g = \tilde{h}_2 \gamma h_2^{-1} = \left(\begin{array}{c|c} \frac{A - xC}{C} & \frac{-Ax + B + xCx - x {}^t A}{-Cx + {}^t A} \end{array} \right) = \left(\begin{array}{cc|cc} 0 & 0 & -C_1^{-1} & 0 \\ 0 & A_4 & 0 & B_4 \\ \hline C_1 & 0 & 0 & 0 \\ 0 & 0 & 0 & {}^t A_4 \end{array} \right).$$

Since $g\tilde{g} = I$ we may apply Lemma 3.8 and Lemma 3.10 to find $Z_r \in \mathfrak{h}_r$ and $Z_{n-r} \in \mathfrak{h}_{n-r}$ so that

$$\begin{pmatrix} 0 & -C_1^{-1} \\ C_1 & 0 \end{pmatrix} Z_r = -\bar{Z}_r \text{ and } \begin{pmatrix} A_4 & B_4 \\ 0 & {}^t A_4 \end{pmatrix} Z_{n-r} = -\bar{Z}_{n-r}.$$

Set $Z = \begin{pmatrix} Z_r & 0 \\ 0 & Z_{n-r} \end{pmatrix}$. Then $gZ = -\bar{Z}$ hence the corresponding cocycle is cohomologically trivial. $\square$

3.13. Proposition. *Let $\gamma \in \mathbf{Sp}(2n, \mathbb{R})$ and suppose that $\tilde{\gamma}\gamma$ is contained in a torsion-free subgroup of $\mathbf{Sp}(2n, \mathbb{R})$. Then the following three statements are equivalent:*

1. $\mathfrak{h}_n^\gamma \neq \phi$
2. f_γ is a cocycle (i.e., $\tilde{\gamma}\gamma = I$).
3. There exists $g \in \mathbf{Sp}(2n, \mathbb{R})$ such that $\gamma = \tilde{g}g^{-1}$.

In this case $\mathfrak{h}_n^\gamma = gC_n$. Moreover such an element g is uniquely determined up to multiplication on the right by an element of $\mathbf{GL}(n, \mathbb{R})$. If $\Gamma \subset \mathbf{Sp}(2n, \mathbb{R})$ is a torsion free arithmetic group and if $\gamma_1, \gamma_2 \in \Gamma$ are distinct then $\mathfrak{h}_n^{\gamma_1} \cap \mathfrak{h}_n^{\gamma_2} = \phi$.

3.14. Proof. Suppose $Z \in \mathfrak{h}_n^\gamma \neq \phi$. Then $\tilde{\gamma}\gamma Z = \tilde{\gamma}\tilde{Z} = Z$ so $\tilde{\gamma}\gamma$ is torsion. By hypothesis this implies that $\tilde{\gamma}\gamma = I$, so (1) implies (2). If f_γ is a 1-cocycle then by Proposition 3.2 it is a coboundary. Then Lemma 3.5 provides $g \in \mathbf{Sp}(2n, \mathbb{R})$ so that $\gamma = \tilde{g}g^{-1}$, which shows that (2) implies (3). It is easy to see that (3) implies (1).

If $\gamma = \tilde{g}g^{-1} = \tilde{g}'(g')^{-1}$ then τ fixes $\alpha = (g')^{-1}g$. Hence $\alpha \in \mathbf{GL}(n, \mathbb{R})$, which proves the next statement.

Finally, suppose $\gamma_1, \gamma_2 \in \Gamma$ which is torsion-free. If there exists $Z \in \mathfrak{h}_n^{\gamma_1} \cap \mathfrak{h}_n^{\gamma_2}$ then $\gamma_1 Z = \tau(Z) = \gamma_2 Z$ so $\gamma_2^{-1}\gamma_1$ fixes Z , hence $\gamma_1 = \gamma_2$. $\square$

4. THE VARIETY $\Gamma(4m) \backslash \mathfrak{h}_n$

4.1. In this section we will establish a strengthening of Proposition 3.13 which applies to (certain) arithmetic subgroups of $\mathbf{Sp}(2n, \mathbb{R})$. We shall be concerned with the following arithmetic groups:

- $\Gamma(1) = \mathbf{Sp}(2n, \mathbb{Z})$
- $\Gamma(N) = \{\gamma \in \Gamma(1) \mid \gamma \equiv I \pmod{N}\}$
- $\Gamma_{2m}(2) = \left\{ \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \Gamma \mid \begin{array}{l} A, D \equiv I \pmod{2} \\ B, C \equiv 0 \pmod{2m} \end{array} \right\}$
- $\Gamma_\ell(1) = \mathbf{GL}(n, \mathbb{Z})$
- $\Gamma_\ell(N) = \{\gamma \in \Gamma_\ell(1) \mid \gamma \equiv I \pmod{N}\}$

(The analogous construction of $\Gamma_m(2)$ does not yield a group unless m is even.) Throughout the remainder of the paper we will be interested in the case $N = 4m$ and $m \geq 1$. The following lemma (which depends on Proposition 4.8 below) is not essential for this paper however it describes the relations between the above groups.

4.2. Lemma. *The short exact sequence*

$$1 \rightarrow \Gamma(4m) \rightarrow \Gamma_{2m}(2) \rightarrow \Gamma(4m) \backslash \Gamma_{2m}(2) \rightarrow 1 \quad (4.2.1)$$

induces a bijection

$$H^1(\mathbb{C}/\mathbb{R}, \Gamma(4m)) \cong \Gamma(4m) \backslash \Gamma_{2m}(2) / \Gamma_\ell(2).$$

4.3. Proof. We claim that $H^1(\mathbb{C}/\mathbb{R}, \Gamma(4m)) \rightarrow H^1(\mathbb{C}/\mathbb{R}, \Gamma_{2m}(2))$ is trivial. For if $\gamma \in \Gamma(4m)$ and if f_γ is a 1-cocycle then by Proposition 3.13 (below), $\mathfrak{h}_n^\gamma \neq \phi$. Then Proposition 4.8 implies that $\gamma = \tilde{g}g^{-1}$ for some $g \in \Gamma_{2m}(2)$, which proves the claim. Next, observe that $\text{Gal}(\mathbb{C}/\mathbb{R})$ acts trivially on the quotient group $\Gamma(4m) \backslash \Gamma_{2m}(2)$. (For if $g \in \Gamma_{2m}(2)$ then $g - \tilde{g} \equiv 0 \pmod{4m}$ so $\tilde{g}g^{-1} \equiv I \pmod{4m}$.) Therefore the long exact cohomology sequence associated to (4.2.1) is:

$$\begin{array}{ccccc} H^0(\Gamma(4m)) & \rightarrow & H^0(\Gamma_{2m}(2)) & \rightarrow & H^0(\Gamma(4m) \backslash \Gamma_{2m}(2)) \rightarrow H^1(\Gamma(4m)) \rightarrow 1 \\ \parallel & & \parallel & & \parallel \\ \Gamma_\ell(4m) & & \Gamma_\ell(2) & & \Gamma(4m) \backslash \Gamma_{2m}(2) \end{array}$$

which completes the proof of Lemma 4.2. $\square$

Let $X = \Gamma(N) \backslash \mathfrak{h}_n$. Then X has the structure of a quasi-projective complex algebraic variety. The involution $\tau(Z) = -\overline{Z}$ passes to an anti-holomorphic involution on X and defines a real structure (let us call it the τ -real structure) on X . In this section we will determine the real points of X . (It is a theorem of Shimura [Sh] that the Baily-Borel compactification V of X even admits the structure of an algebraic variety defined over the rational numbers. See §8.) Define

$$S = \bigcup_{\gamma \in \Gamma(4m)} \mathfrak{h}_n^\gamma \subset \mathfrak{h}_n$$

to be the set of all $\Gamma(4m)$ -real points in $\mathfrak{h}_n$. Denote by $\Gamma(4m) \backslash S$ the image of S in $X = \Gamma(4m) \backslash \mathfrak{h}_n$. For $g \in \Gamma_{2m}(2)$ set ${}^g\Gamma_\ell(4m) = g\Gamma_\ell(4m)g^{-1}$. In this section we will prove the following

4.4. Theorem. *The set $X_{\mathbb{R}}$ of real points of X is precisely $\Gamma(4m) \backslash S$. It consists of the disjoint union*

$$X_{\mathbb{R}} = \coprod_g {}^g\Gamma_{\ell}(4m) \backslash giC_n \quad (4.4.1)$$

of finitely many copies of $\Gamma_{\ell}(4m) \backslash C_n$, indexed by elements

$$g \in \Gamma(4m) \backslash \Gamma_{2m}(2) / \Gamma_{\ell}(2) = H^1(\mathbb{C}/\mathbb{R}, \Gamma(4m)). \quad (4.4.2)$$

The copy indexed by $g \in \Gamma_{2m}(2)$ corresponds to the cohomology class of f_{γ} where $\gamma = \tilde{g}g^{-1} \in \Gamma(4m)$.

In order to identify the set of real points $X_{\mathbb{R}}$ we will make repeated use of the following observation: *the image $[Z] \in X$ of a point $Z \in \mathfrak{h}_n$ lies in $X_{\mathbb{R}}$ iff there exists $\gamma \in \Gamma(4m)$ such that $\gamma Z = -\bar{Z}$ (i.e. such that $Z \in \mathfrak{h}_n^{\gamma}$).* The following lemma will be used throughout the next few sections, and it also accounts for the indexing set in (4.4.2).

4.5. Lemma. *Let $g \in \mathbf{Sp}(2n, \mathbb{Z})$. Fix $m \geq 1$.*

1. *If $g \in \Gamma(m)$ then $\tilde{g}g^{-1} \in \Gamma(2m)$.*
2. *If $g \in \Gamma(2m)$ then $\tilde{g}g \in \Gamma(4m)$.*
3. *If $\tilde{g}g^{-1} \in \Gamma(4m)$ then $g = \beta u$ for some $\beta \in \Gamma_{2m}(2)$ and for some $u \in \mathbf{GL}(n, \mathbb{Z})$.*

4.6. Proof. Let $g = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$. Since $\tilde{g} = g - \begin{pmatrix} 0 & 2B \\ 2C & 0 \end{pmatrix}$ we have

$$\tilde{g}g^{-1} = I - \begin{pmatrix} -2B^tC & 2B^tA \\ 2C^tD & -2C^tA \end{pmatrix}. \quad (4.6.1)$$

This gives part (1). If $g = I + 2mg'$ then $\tilde{g} + g$ is even, and $\tilde{g}g = I + 2m(\tilde{g}' + g') + 4m^2\tilde{g}'g$ which proves part(2). Now suppose that $\tilde{g}g^{-1} \in \Gamma(4m)$ so that $B^tC \equiv B^tA \equiv C^tD \equiv 0 \pmod{2m}$. Then

$$\begin{aligned} B &= B({}^tAD - {}^tCB) = (B^tA)D - (B^tC)B \equiv 0 \pmod{2m} \\ {}^tC &= ({}^tAD - {}^tCB){}^tC = {}^tA(D^tC) - {}^tC(B^tC) \equiv 0 \pmod{2m}. \end{aligned}$$

Moreover, since $C^tB \equiv 0 \pmod{2m}$ and $A^tD - B^tC = I$ we see that $A^tD \equiv I \pmod{2}$. Hence A is invertible $\pmod{2}$. But reduction $\pmod{2}$ is a surjective mapping $\mathbf{GL}(n, \mathbb{Z}) \rightarrow \mathbf{GL}(n, \mathbb{Z}/(2))$ hence there exists $U \in \mathbf{GL}(n, \mathbb{Z})$ so that $U \equiv A \pmod{2}$ from which it also follows that $U^{-1} \equiv {}^tD \pmod{2}$. Let $u = \begin{pmatrix} U & 0 \\ 0 & {}^tU^{-1} \end{pmatrix} \in \mathbf{Sp}(2n, \mathbb{R})$. Then

$$\beta = gu^{-1} = \begin{pmatrix} AU^{-1} & B^tU \\ CU^{-1} & D^tU \end{pmatrix}$$

is in $\Gamma_{2m}(2)$ as claimed. Conversely, if $\beta \in \Gamma_{2m}(2)$ and $u \in \mathbf{GL}(n, \mathbb{Z})$ then $\tau(\beta u)(\beta u)^{-1} = \tilde{\beta}\beta^{-1}$ which is easily seen (using 4.6.1) to lie in $\Gamma(4m)$. $\square$

The following proposition is a consequence of the theorem of Silhol [Si] (Thm. 1.4) and Comessatti, and will be proven in §7.

4.7. Proposition. *Let $\gamma \in \Gamma(2)$ and suppose that $Z \in \mathfrak{h}_n$ is not fixed by any element of $\mathbf{Sp}(2n, \mathbb{Z})$ other than $\pm I$. Suppose that $\tilde{Z} = \gamma Z$. Then there exists $h \in \mathbf{Sp}(2n, \mathbb{Z})$ such that $\gamma = \tilde{h}h^{-1}$, hence $\mathfrak{h}_n^\gamma = h \cdot iC_n$.*

4.8. Corollary. *Suppose $\gamma \in \Gamma(4m)$ and $\mathfrak{h}_n^\gamma \neq \emptyset$. Then there exists $g \in \Gamma_{2m}(2)$ such that $\gamma = \tilde{g}g^{-1}$ (hence $\mathfrak{h}_n^\gamma = giC_n$).*

4.9. Proof. Let $Z \in \mathfrak{h}_n^\gamma$ be a point which is not fixed by any $g \in \mathbf{Sp}(2n, \mathbb{Z})$ other than $\pm I$. Such points exist, and even form a dense subset of $\mathfrak{h}_n^\gamma$ because by Proposition 3.13, the set $\mathfrak{h}_n^\gamma$ is a translate (by some element of $\mathbf{Sp}(2n, \mathbb{R})$) of the cone iC_n . By Proposition 4.7, there exists $h \in \mathbf{Sp}(2n, \mathbb{Z})$ so that $\gamma = \tilde{h}h^{-1}$. By Lemma 4.5 this implies that $h = \beta u$ for some $\beta \in \Gamma_{2m}(2)$ and some $u \in \mathbf{GL}(n, \mathbb{Z})$. Hence $\gamma = \tilde{h}h^{-1} = \tilde{\beta}\beta^{-1}$ as claimed. $\square$

4.10. Lemma. *Let $g_1, g_2 \in \Gamma_{2m}(2)$. Suppose there exist $Z_1 \in g_1iC_n$, $Z_2 \in g_2iC_n$, and $\gamma \in \Gamma(4m)$ such that $\gamma Z_1 = Z_2$. Then there exist $\alpha \in \Gamma_\ell(2)$ such that $g_2 = \gamma g_1 \alpha$.*

4.11. Proof. By section 2.2 (3) we have a commutative diagram

$$\begin{array}{ccc} Z_1 & \xrightarrow{\gamma} & Z_2 \\ \tilde{g}_1 g_1^{-1} \downarrow & & \downarrow \tilde{g}_2 g_2^{-1} \\ \tilde{Z}_1 & \xrightarrow{\tilde{\gamma}} & \tilde{Z}_2 \end{array}$$

so the element $\theta = g_1 \tilde{g}_1^{-1} \tilde{\gamma}^{-1} \tilde{g}_2 g_2^{-1} \gamma$ fixes Z_1 . By lemma 4.5, $\theta \in \Gamma(4m)$ which is torsion-free. So $\theta = I$, which implies

$$\gamma = g_2 \tilde{g}_2^{-1} \tilde{\gamma} \tilde{g}_1 g_1^{-1}$$

Therefore, $\tau(g_2^{-1} \gamma g_1) = g_2^{-1} \gamma g_1$ which implies that $g_2^{-1} \gamma g_1 \in \mathbf{GL}_n(\mathbb{R}) \cap \Gamma_{2m}(2) = \Gamma_\ell(2)$. $\square$

4.12. Proof of Theorem 4.4. For $\gamma \in \Gamma(4m)$ define $X^\gamma \subset X = \Gamma(4m) \backslash \mathfrak{h}_n$ to be the image of $\mathfrak{h}_n^\gamma$ in X . For each $g \in \Gamma_{2m}(2)$ the corresponding element $\gamma = \tau(g)g^{-1}$ lies in $\Gamma(4m)$, and $\mathfrak{h}_n^\gamma = giC_n$. (This follows from lemma 4.5 and Proposition 3.13.) The resulting mapping

$$\bigcup_{g \in \Gamma_{2m}(2)} giC_n \longrightarrow \bigcup_{\gamma \in \Gamma(4m)} \mathfrak{h}_n^\gamma$$

is surjective by Corollary 4.8. For each $g \in \Gamma_{2m}(2)$ the composition $giC_n \rightarrow \mathfrak{h}_n^\gamma \rightarrow X^\gamma$ is surjective and induces an isomorphism ${}^g\Gamma_\ell(4m) \backslash giC_n \rightarrow X^\gamma$ (where ${}^g\Gamma_\ell(4m) = g\Gamma_\ell(4m)g^{-1}$.) By lemma 4.10 this determines a bijection

$$\coprod_{g \in \Gamma(4m) \backslash \Gamma_{2m}(2) / \Gamma_\ell(2)} {}^g\Gamma_\ell(4m) \backslash giC_n \longrightarrow \Gamma(4m) \backslash \bigcup_{\gamma \in \Gamma(4m)} \mathfrak{h}_n^\gamma = X_\mathbb{R}$$

5. UPPER HALF-PLANE

For the upper half-plane $\mathfrak{h}_1$ it is possible to do a little better since the group $\Gamma(2)/\pm I$ acts freely.

5.1. Lemma. *A point $Z \in \mathfrak{h}_1$ is $(\Gamma(2), \tau)$ -real if and only if gZ is $(\Gamma(2), \tau)$ -real, for every $g \in SL(2, \mathbb{Z})$.*

5.2. Proof. Suppose $\tau(Z) = \gamma Z$ for some $\gamma \in \Gamma(2)$. Then

$$\tau(gz) = \tau(g)\tau(z) = (\tau(g)g^{-1})(g\gamma g^{-1})gZ \in \Gamma(2) \cdot \Gamma(2) \cdot gZ. \quad \square$$

5.3. Lemma. *If a point $y \in \mathfrak{h}_1$ is $(\Gamma(2), \tau)$ -real, then there exists $\beta \in SL(2, \mathbb{Z})$ and there exists $Y \in C_1 = \mathbb{R}_+$ so that $y = \beta \cdot iY$.*

5.4. Proof. Say $\tilde{y} = \gamma y$ for some $\gamma \in \Gamma(2)$. We consider two cases: (a) when y is not fixed by any element of $SL_2(\mathbb{Z})$ other than $\pm I$ and (b) when y is fixed by some other nontrivial element of $SL_2(\mathbb{Z})$.

In case (a), the result is just Proposition 4.7. Now consider case (b). There are two classes of elements which are fixed by nontrivial subgroups of $SL_2(\mathbb{Z})$. These are the translates (by elements of $SL_2(\mathbb{Z})$) of i and the translates of $\omega = e^{2\pi i/3}$. In the first case, $y = \beta \cdot i$ (for some $\beta \in SL_2(\mathbb{Z})$) and the conclusion follows. In the second case, the stabilizer of ω is the subgroup

$$S = \left\{ \pm I, \pm \begin{pmatrix} 0 & -1 \\ 1 & 1 \end{pmatrix}, \pm \begin{pmatrix} -1 & -1 \\ 1 & 0 \end{pmatrix} \right\}.$$

Suppose $y = g\omega$ (for some $g \in SL_2(\mathbb{Z})$). We claim that y is not $\Gamma(2)$ -real. Suppose otherwise. Then $\tilde{y} = \gamma y$ gives $\tilde{\omega} = \gamma'\omega$ where

$$\gamma' = \tilde{g}^{-1}\gamma g = (\tilde{g}^{-1}g)g^{-1}\gamma g \in \Gamma(2)$$

by Lemma 4.5. But $\tilde{\omega} = -\bar{\omega} = j\omega$ where $j = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$. So $j^{-1}\gamma'$ stabilizes ω or $\gamma' = js$ for some $s \in S$. However none of these matrices js lies in $\Gamma(2)$, which is a contradiction. This completes the proof of Lemma 5.3. $\square$

The group $\Gamma(2)$ does not induce any identifications on the cone iC_1 . However iC_1 is preserved by the subgroup $S' = \{\pm I, \pm v\}$ where $v = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$. So we obtain

5.5. Theorem. *The (τ) -real points of $X = \Gamma(2) \backslash \mathfrak{h}_1$ consists of the union of 3 copies of the cone $C_1 \cong \mathbb{R}_+$,*

$$X_{\mathbb{R}} = \bigcup_{\beta \in \Gamma(2) \backslash SL_2(\mathbb{Z}) / S'} \beta \cdot iC_1. \quad \square$$

We remark that in the Baily-Borel compactification $V \cong \mathbb{P}^1$ these three half-lines are glued end-to-end to form a single circle.

6. MODULI SPACE INTERPRETATION

6.1. Recall [La] that a symplectic form Q on $\mathbb{C}^n$ is *compatible* with the complex structure if $Q(iu, iv) = Q(u, v)$ for all $u, v \in \mathbb{C}^n$. A compatible form Q is *positive* if the symmetric form $R(u, v) = Q(iu, v)$ is positive definite. If Q is compatible and positive then it is the imaginary part of a unique positive definite Hermitian form $H = R + iQ$. Let $L \subset \mathbb{C}^n$ be a lattice and let $H = R + iQ$ be a positive definite Hermitian form on $\mathbb{C}^n$. A basis of L is *symplectic* if the matrix for Q with respect to this basis is $\begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix}$. The lattice L is symplectic if it admits a symplectic basis.

A principally polarized abelian variety is a pair $(A = \mathbb{C}^n/L, H = R + iQ)$ where H is a positive definite Hermitian form on $\mathbb{C}^n$ and where $L \subset \mathbb{C}^n$ is a symplectic lattice relative to $Q = \text{Im}(H)$. A real structure on (A, H) is a complex anti-linear involution $\kappa : \mathbb{C}^n \rightarrow \mathbb{C}^n$ such that $\kappa(L) = L$. A real structure κ on (A, H) is compatible with the polarization H if $Q(\kappa u, \kappa v) = -Q(u, v)$ for all $u, v \in \mathbb{C}^n$. In this case (following [Si]) we refer to the triple (A, H, κ) as a *real principally polarized abelian variety*. If (A', H', κ') is another such, then an isomorphism between them is a complex linear mapping $\phi : \mathbb{C}^n \rightarrow \mathbb{C}^n$ such that

- (a.) $\phi(L) = L'$,
- (b.) $\phi_*(H) = H'$,
- (c.) $\phi_*(\kappa) = \kappa'$, that is, $\phi\kappa\phi^{-1} = \kappa'$.

Let $(A = \mathbb{C}^n/L, H = R + iQ)$ be a principally polarized abelian variety. A level N structure on A is a choice of basis $\{U_i, V_j\}$ (with $1 \leq i, j \leq n$) for the N -torsion points of A , which is symplectic, in the sense that there exists a symplectic basis $\{u_i, v_j\}$ for L such that

$$U_i \equiv \frac{u_i}{N} \text{ and } V_j \equiv \frac{v_j}{N} \pmod{L}$$

(for $1 \leq i, j \leq n$). For a given level N structure, such a choice $\{u_i, v_j\}$ determines a mapping

$$F : \mathbb{R}^n \oplus \mathbb{R}^n \rightarrow \mathbb{C}^n \tag{6.1.1}$$

such that $F(\mathbb{Z}^n \oplus \mathbb{Z}^n) = L$, by $F(e_i) = u_i$ and $F(f_j) = v_j$ where $\{e_i, f_j\}$ (with $1 \leq i, j \leq n$) is the standard basis of $\mathbb{R}^n \oplus \mathbb{R}^n$. The choice $\{u_i, v_j\}$ (or equivalently, the mapping F) will be referred to as a *lift* of the level N structure. It is well defined modulo the principal congruence subgroup $\Gamma(N)$, that is, if $F' : \mathbb{R}^n \oplus \mathbb{R}^n \rightarrow \mathbb{C}^n$ is another lift of the level structure, then $F' \circ F^{-1} \in \Gamma(N)$.

A level N -structure $\{U_i, V_j\}$ is *compatible* with a real structure κ if for some (and hence for any) lift $\{u_i, v_j\}$ of the level structure,

$$\kappa\left(\frac{u_i}{N}\right) \equiv -\frac{u_i}{N} \pmod{L} \text{ and } \kappa\left(\frac{v_j}{N}\right) \equiv +\frac{v_j}{N} \pmod{L}$$

for all $1 \leq i, j \leq n$. In other words, the following diagram commutes (mod L).

$$\begin{array}{ccc} \frac{1}{N}(\mathbb{Z}^n \oplus \mathbb{Z}^n) & \xrightarrow{F} & \frac{1}{N}L \\ \downarrow \iota_- & & \downarrow \kappa \\ \frac{1}{N}(\mathbb{Z}^n \oplus \mathbb{Z}^n) & \xrightarrow{F} & \frac{1}{N}L \end{array} \quad (6.1.2)$$

where (cf. §2.2) $\iota_- = \begin{pmatrix} -I & 0 \\ 0 & I \end{pmatrix}$

6.2. Definition. A real principally polarized abelian variety with level N structure is a quadruple $\mathcal{A} = (A = \mathbb{C}^n/L, H = R + iQ, \kappa, \{U_i, V_j\})$ where (A, H, κ) is a real principally polarized abelian variety and where $\{U_i, V_j\}$ is a level N -structure which is compatible with κ . An isomorphism

$$\mathcal{A} = (A, H, \kappa, \{U_i, V_j\}) \cong (A', H', \kappa', \{U'_i, V'_j\}) = \mathcal{A}' \quad (6.2.1)$$

is a complex linear mapping $\phi : \mathbb{C}^n \rightarrow \mathbb{C}^n$ such that (a.), (b.), and (c.) of §6.1 hold, and such that

$$(d.) \quad \phi\left(\frac{u_i}{N}\right) \equiv \frac{u'_i}{N} \pmod{L'} \text{ and } \phi\left(\frac{v_j}{N}\right) \equiv \frac{v'_j}{N} \pmod{L'} \text{ for } 1 \leq i, j \leq n$$

for some (and hence for any) lift $\{u_i, v_j\}$ and $\{u'_i, v'_j\}$ of the level structures.

6.3. Fix $N \geq 1$. Each $Z \in \mathfrak{h}_n$ determines a principally polarized abelian variety (A_Z, H_Z) with level N structure as follows. Let Q_0 be the standard symplectic form on $\mathbb{R}^n \oplus \mathbb{R}^n$ with matrix $\begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix}$ (with respect to the standard basis of $\mathbb{R}^n \oplus \mathbb{R}^n$). Let $F_Z : \mathbb{R}^n \oplus \mathbb{R}^n \rightarrow \mathbb{C}^n$ be the real linear mapping with matrix (Z, I) , that is,

$$F_Z \begin{pmatrix} x \\ y \end{pmatrix} = Zx + y.$$

Then $Q_Z = (F_Z)_*(Q_0)$ is a compatible, positive symplectic form and $L_Z = F_Z(\mathbb{Z}^n \oplus \mathbb{Z}^n)$ is a symplectic lattice with symplectic basis $F_Z(\text{standard basis})$. The Hermitian form corresponding to Q_Z is

$$H_Z(u, v) = Q_Z(iu, v) + iQ_Z(u, v) = {}^t u(\text{Im}(Z))^{-1} \bar{v} \text{ for } u, v \in \mathbb{C}^n.$$

Then the pair $(A_Z = \mathbb{C}^n/L_Z, H_Z)$ is a principally polarized abelian variety. If z_1, z_2 are the standard coordinates on $\mathbb{C}^n$ then, with respect to the above symplectic basis of L , the differential forms dz_1, dz_2 have period matrix (Z, I) . If $\{e_1, \dots, e_n, f_1, \dots, f_n\}$ denote the standard basis elements of $\mathbb{R}^n \oplus \mathbb{R}^n$ then the collection $\{F_Z(e_i/N), F_Z(f_i/N)\} \pmod{L}$ is a level N structure on (A_Z, H_Z) , which we refer to as the *standard level N structure*.

Let $Z, \Omega \in \mathfrak{h}_n$. Suppose $\psi : (A_\Omega = \mathbb{C}^n/L_\Omega, H_\Omega) \rightarrow (A_Z = \mathbb{C}^n/L_Z, H_Z)$ is an isomorphism of the corresponding principally polarized abelian varieties, that is, $\psi(L_\Omega) = L_Z$ and $\psi_*(H_\Omega) = H_Z$. Set $h = {}^t(F_Z^{-1}\psi F_\Omega) = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$. Then: $h \in \mathbf{Sp}(2n, \mathbb{Z})$, $\Omega = h \cdot Z$, and $\psi(M) = {}^t(CZ + D)M$ for all $M \in \mathbb{C}^n$. Since $h \cdot Z$ is symmetric, the following diagram

commutes:

$$\begin{array}{ccccccc}
\begin{pmatrix} x \\ y \end{pmatrix} & \mathbb{R}^n \oplus \mathbb{R}^n & \xrightarrow{F_{h \cdot Z}} & \mathbb{C}^n & & M & \\
\downarrow & \downarrow & & \downarrow \psi & & \downarrow & \\
{}^t h \begin{pmatrix} x \\ y \end{pmatrix} & \mathbb{R}^n \oplus \mathbb{R}^n & \xrightarrow{F_Z} & \mathbb{C}^n & & {}^t(CZ + D)M &
\end{array} \tag{6.3.1}$$

Let $Z \in \mathfrak{h}_n$ with $\tilde{Z} = \tau(Z) = -\overline{Z}$. Then the following diagram commutes:

$$\begin{array}{ccccccc}
\begin{pmatrix} x \\ y \end{pmatrix} & \mathbb{R}^n \oplus \mathbb{R}^n & \xrightarrow{F_Z} & \mathbb{C}^n & & M & \\
\downarrow & \downarrow & & \downarrow & & \downarrow & \\
{}^t I_- \begin{pmatrix} x \\ y \end{pmatrix} & \mathbb{R}^n \oplus \mathbb{R}^n & \xrightarrow{F_{\tau(Z)}} & \mathbb{C}^n & & \overline{M} &
\end{array} \tag{6.3.2}$$

6.4. The points in $\mathfrak{h}_n^{\Gamma(1)}$ give rise to abelian varieties with a real structure, as follows. Suppose $Z \in \mathfrak{h}_n$, $\gamma = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \mathbf{Sp}(2n, \mathbb{Z})$ and suppose that $\gamma \cdot Z = \tau(Z)$. Define

$$\kappa(\gamma, Z) : \mathbb{C}^n \rightarrow \mathbb{C}^n \text{ by } M \mapsto {}^t(CZ + D)\overline{M}. \tag{6.4.1}$$

Then $\kappa(\gamma, Z)$ is a real structure on (A_Z, H_Z) which is compatible with the polarization H_Z , and the following diagram (which is the composition of the two preceding diagrams) commutes:

$$\begin{array}{ccccccc}
\begin{pmatrix} x \\ y \end{pmatrix} & \mathbb{R}^n \oplus \mathbb{R}^n & \xrightarrow{F_Z} & \mathbb{C}^n & & M & \\
\downarrow & \downarrow & & \downarrow & & \downarrow \kappa(\gamma, Z) & \\
{}^t \gamma {}^t I_- \begin{pmatrix} x \\ y \end{pmatrix} & \mathbb{R}^n \oplus \mathbb{R}^n & \xrightarrow{F_Z} & \mathbb{C}^n & & {}^t(CZ + D)\overline{M} &
\end{array} \tag{6.4.2}$$

6.5. Proposition. *Let $Z \in \mathfrak{h}_n$ and $\gamma \in \mathbf{Sp}(2n, \mathbb{R})$ and suppose that $\tilde{Z} = \gamma \cdot Z$. Then $\gamma \in \Gamma(N)$ iff the real structure $\kappa(\gamma, Z)$ on (A_Z, H_Z) is compatible with the standard level N structure, and in this case the quadruple*

$$(A_Z, H_Z, \kappa(\gamma, Z), \{F_Z(e_i/N), F_Z(f_j/N)\})$$

is a real principally polarized abelian variety with (compatible) level N structure.

The proof follows immediately from the diagrams (6.4.2) and (6.1.2). We remark that if $N \geq 3$, and if $Z \in \mathfrak{h}_n^{\Gamma(N)}$ then there is a unique $\gamma \in \Gamma(N)$ such that $\tilde{Z} = \gamma \cdot Z$ so we may unambiguously denote $\kappa(\gamma, Z)$ simply by κ_Z . In this case, denote by $\mathcal{A}_Z$ the resulting real abelian variety with principal polarization and (compatible) level N structure. The proof of the following theorem will appear in §7.

6.6. Theorem. Fix $N = 4m$ with $m \geq 1$. The association $Z \mapsto \mathcal{A}_Z$ determines a one to one correspondence between the points of $X_{\mathbb{R}}$ and the set of isomorphism classes of real principally polarized abelian varieties with level N structure.

7. THE COMESSATTI LEMMA

Recall the following theorem of Silhol [Si] Thm. 1.4., (also [Si2]) and Comessatti [C].

7.1. Theorem. Let (A, H, κ) be a real principally polarized abelian variety. Then there exists $Z = X + iY \in \mathfrak{h}_n$ so that $2X$ is integral, and there exists an isomorphism of real principally polarized abelian varieties,

$$(A, H, \kappa) \cong (A_Z, H_Z, \sigma)$$

where $\sigma(M) = \overline{M}$ is complex conjugation.

For the purposes of this paper we will need a slight restatement of this result.

7.2. Lemma. (A) Let $Z \in \mathfrak{h}_n$ and suppose that $\tilde{Z} = \gamma \cdot Z$ for some $\gamma \in \mathbf{Sp}(2n, \mathbb{Z})$. Then Z is equivalent under $\mathbf{Sp}(2n, \mathbb{Z})$ to an element $X + iY \in \mathfrak{h}_n$ such that $2X$ is integral.

(B) Let (A, H, κ) be a real principally polarized abelian variety. Then there exists $\gamma \in \mathbf{Sp}(2n, \mathbb{Z})$ and there exists $Z = X + iY \in \mathfrak{h}_n$ such that $\tilde{Z} = \gamma \cdot Z$ and $2X$ is integral; and there exists an isomorphism of real principally polarized abelian varieties

$$(A, H, \kappa) \cong (A_Z, H_Z, \kappa(\gamma, Z)).$$

7.3. Proof. For part (B), Theorem 7.1 provides $Z = X + iY \in \mathfrak{h}_n$ so that $2X$ is integral. Take $\gamma = \begin{pmatrix} I & -2X \\ 0 & I \end{pmatrix}$. Then $\tilde{Z} = -\overline{Z} = \gamma \cdot Z$ and $\kappa(\gamma, Z)(M) = \overline{M} = \sigma(M)$ by (6.4.2).

For part (A), given Z and γ , apply part (B) to the real abelian variety $(A_Z, H_Z, \kappa(\gamma, Z))$ to obtain an isomorphic real abelian variety $(A_{Z'}, H_{Z'}, \kappa(\gamma', Z'))$ such that $Z' = X' + iY'$ with $2X'$ integral. Then (A_Z, H_Z) and $(A_{Z'}, H_{Z'})$ are isomorphic principally polarized abelian varieties so there exists $g \in \mathbf{Sp}(2n, \mathbb{Z})$ with $Z' = g \cdot Z$. $\square$

7.4. Proof of Proposition 4.7. By Comessatti's lemma, Z is equivalent (via some $h \in \mathbf{Sp}(2n, \mathbb{Z})$) to some element $X + iY \in \mathfrak{h}_n$ with $2X \in M_{n \times n}(\mathbb{Z})$. We claim, in this case, that X is integral. Translation by X is given by the symplectic matrix

$$T_X = \begin{pmatrix} I & X \\ 0 & I \end{pmatrix}$$

so we may write $Z = h(X + iY) = hT_X iY$. Then $\tau Z = \tau(hT_X) iY = \gamma Z$, so the following element $\gamma^{-1} \tau(hT_X) (hT_X)^{-1}$ fixes Z . By our assumption on Z , this implies that

$$\gamma = \pm I \tau(hT_X) (hT_X)^{-1} \tag{7.4.1}$$

or,

$$\pm T_{-2X} = \tilde{h}^{-1} \gamma h = (\tilde{h}^{-1} h) (h^{-1} \gamma h) \in \Gamma(2). \Gamma(4m)$$

(using Lemma 4.5 and the fact that $\Gamma(4m)$ is normal in $\mathbf{Sp}(2n, \mathbb{Z})$). So $2X$ is “even”, hence X is integral.

If the plus sign occurs in (7.4.1) then $\gamma = \tau(hT_X)(hT_X)^{-1}$ hence $\mathfrak{h}_n^\gamma = hT_X \cdot iC_n$. If the minus sign occurs, set $\omega = \begin{pmatrix} 0 & -I \\ I & 0 \end{pmatrix}$. Then $\gamma = \tau(hT_X\omega)(hT_X\omega)^{-1}$ and hence $\mathfrak{h}_n^\gamma = hT_X\omega \cdot iC_n$. $\square$

In the next proposition we strengthen these results to include level structures.

7.5. Proposition. *Suppose $N \geq 3$. Suppose $\mathcal{A}$ is a real abelian variety with principal polarization and level N structure. Then there exists $Z \in \mathfrak{h}_n$, there exists $\gamma \in \Gamma(N)$ such that $\gamma \cdot Z = \tau(Z)$, and there exists an isomorphism*

$$\phi : \mathcal{A}_Z \rightarrow \mathcal{A}$$

of real principally polarized abelian varieties with level structures. If $N = 4m$ (and $m \geq 1$) the cohomology class $[f_\gamma] \in H^1(\mathbb{C}/\mathbb{R}, \Gamma(4m))$ is uniquely determined by the isomorphism class of $\mathcal{A}$.

7.6. Proof. Write $\mathcal{A} = (A = \mathbb{C}^n/L, H = R + iQ, \kappa, \{U_i, V_j\})$. By Lemma 7.2, there exists $Z' \in \mathfrak{h}_n$, there exists $\gamma' \in \mathbf{Sp}(2n, \mathbb{Z})$ with $\gamma' \cdot Z' = \tau(Z')$, and there exists $\phi' : \mathbb{C}^n \rightarrow \mathbb{C}^n$ such that ϕ' induces an isomorphism $(A, H, \kappa) \cong (A_{Z'}, H_{Z'}, \kappa(\gamma', Z'))$, that is:

- (a.) $\phi'(L_{Z'}) = L$
- (b.) $\phi'_*(H_{Z'}) = H$
- (c.) $\phi'_*(\kappa(\gamma', Z')) = \kappa$.

The isomorphism ϕ' must be modified because it does not necessarily take the standard level N structure on $(A_{Z'}, H_{Z'})$ to the given level N structure on (A, H) . Choose a lift $\{u_i, v_j\}$ ($1 \leq i, j \leq n$) of the level N structure on (A, H) and let $F : \mathbb{R}^2 \oplus \mathbb{R}^2 \rightarrow \mathbb{C}^2$ be the resulting mapping (6.1.1). Define

$${}^t g^{-1} = F^{-1} \circ \phi' \circ F_{Z'} \in \mathbf{Sp}(2n, \mathbb{Z}) \quad (7.6.1)$$

$$Z = g \cdot Z' \quad (7.6.2)$$

$$\gamma = \tau(g)\gamma'g^{-1} = I_- g I_- \gamma' g^{-1}. \quad (7.6.3)$$

If $g = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$ define $\psi : \mathbb{C}^n \rightarrow \mathbb{C}^n$ by $\psi(M) = {}^t(CZ + D)M$. We claim that $\gamma \in \Gamma(N)$, that $\tilde{Z} = \gamma \cdot Z$ and that the mapping

$$\phi = \phi' \circ \psi : \mathbb{C}^n \rightarrow \mathbb{C}^n$$

induces an isomorphism $\mathcal{A}_Z \cong \mathcal{A}$ between principally polarized real abelian varieties with (compatible) level N structures.

The matrix g is defined so that the bottom square in the following diagram commutes; by (6.3.1) the top square also commutes. The mapping ϕ is the composition down the

right-hand column.

$$\begin{array}{ccc}
\mathbb{R}^n \oplus \mathbb{R}^n & \xrightarrow{F_Z} & \mathbb{C}^n \\
\downarrow \scriptstyle {}^t g & & \downarrow \scriptstyle \psi \\
\mathbb{R}^n \oplus \mathbb{R}^n & \xrightarrow{F_{Z'}} & \mathbb{C}^n \\
\downarrow \scriptstyle {}^t g^{-1} & & \downarrow \scriptstyle \phi' \\
\mathbb{R}^n \oplus \mathbb{R}^n & \xrightarrow{F} & \mathbb{C}^n
\end{array} \tag{7.6.4}$$

Clearly, $\tilde{Z} = \gamma \cdot Z$, $\phi_*(L_Z) = L$ and $\phi_*(H_Z) = H$. Now let us check that

$$\phi_*(\kappa(\gamma, Z)) = \kappa. \tag{7.6.5}$$

By §7.6 (c.) it suffices to check that $\psi\kappa(\gamma, Z)\psi^{-1} = \kappa(\gamma', Z')$. But this follows from direct calculation using $\kappa(\gamma, Z) = F_Z {}^t \gamma {}^t I_- F_Z^{-1}$, $\kappa(\gamma', Z') = F_{Z'} {}^t \gamma' {}^t I_- F_{Z'}^{-1}$, $\psi = F_{Z'} {}^t g F_Z^{-1}$, and (7.6.3).

By (7.6.4), $F = \phi \circ F_Z$. Therefore ϕ preserves the level structures. Since the given level structure on (A, H) is compatible with κ , equation (7.6.5) guarantees that the standard level N structure on (A_Z, H_Z) is compatible with $\kappa(\gamma, Z)$. It follows from Proposition 6.5 that $\gamma \in \Gamma(N)$. In summary, we have shown that $(A_Z, H_Z, \kappa(\gamma, Z), \{F_Z(e_i/N), F_Z(f_j/N)\})$ is a real principally polarized abelian variety with (compatible) level N structure, and that the isomorphism ϕ preserves both the real structure and the level structure.

Finally, an isomorphism $\phi : \mathcal{A} \rightarrow \mathcal{A}'$ (6.2.1) between real principally polarized abelian varieties with level $N = 4m$ structures determines a cohomological equivalence between the corresponding 1-cocycles as follows. Choose $Z, Z' \in \mathfrak{h}_n$ and $\gamma, \gamma' \in \Gamma(4m)$ for $\mathcal{A}$ and $\mathcal{A}'$ respectively, as above, and set

$$g = {}^t(F_Z^{-1} \circ \phi \circ F_{Z'}).$$

Then $g \in \Gamma(4m)$ by §6.2(d.). Since $\phi\kappa\phi^{-1} = \kappa'$, diagrams 6.3.1 and 6.4.2 give

$${}^t g {}^t \gamma {}^t I_- {}^t g^{-1} = {}^t \gamma' I_-$$

or $\gamma' = \tilde{g}^{-1} \gamma g$. Hence $[f_\gamma] = [f_{\gamma'}]$ in $H^1(\mathbb{C}/\mathbb{R}, \Gamma(4m))$. $\square$

7.7. Proof of Theorem 6.6. Let $\mathcal{I}$ denote the moduli space of isomorphism classes of real abelian varieties with principal polarization and level N structure. Let

$$S = \bigcup_{\gamma \in \Gamma(N)} \mathfrak{h}_n^\gamma \subset \mathfrak{h}_n$$

denote the set of $\Gamma(N)$ -real points in $\mathfrak{h}_n$ and let $\Gamma(N) \backslash S$ denote its image in $X = \Gamma(N) \backslash \mathfrak{h}_n$. Let $\Phi : S \rightarrow \mathcal{I}$ be the association

$$Z \mapsto (A_Z, H_Z, \kappa_Z, \{F_Z(e_i/N), F_Z(f_j/N)\}).$$

Here, $\kappa_Z : \mathbb{C}^n \rightarrow \mathbb{C}^n$ is the real structure given by (6.4.2), that is, $\kappa(M) = {}^t(CZ + D)\overline{M}$ where $\gamma = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \Gamma(N)$ is the unique element such that $\gamma \cdot Z = -\overline{Z}$. Then Proposition 7.5 says that Φ is surjective. If $Z, Z' \in S$ and if $Z' = \gamma \cdot Z$ for some $\gamma \in \Gamma(N)$ then by (6.3.1) the real abelian variety $(A_Z, H_Z, \kappa_Z, \{F_Z(e_i/N), F_Z(f_i/N)\})$ is isomorphic to $(A_{Z'}, H_{Z'}, \kappa_{Z'}, \{F_{Z'}(e_i/N), F_{Z'}(f_i/N)\})$. Therefore Φ passes to a mapping $\Gamma(N) \backslash S \rightarrow \mathcal{I}$.

On the other hand, if $Z, \Omega \in S$ and if there is an isomorphism $\psi : \mathcal{A}_\Omega \rightarrow \mathcal{A}_Z$ then by (6.3.1) there exists $h \in \mathbf{Sp}(2n, \mathbb{Z})$ such that $\Omega = h \cdot Z$. Since the mapping ψ also preserves the level structures, it follows also from (6.3.1) that $h \in \Gamma(N)$. Hence the mapping $\Gamma(N) \backslash S \rightarrow \mathcal{I}$ is also injective. By Theorem 4.4, the quotient $\Gamma(N) \backslash S$ is precisely the variety $X_{\mathbb{R}}$ of real points in X . $\square$

8. BAILY-BOREL COMPACTIFICATION

Let $\overline{\mathfrak{h}}_n$ denote the Satake partial compactification of $\mathfrak{h}_n$ which is obtained by attaching all rational boundary components, with the Satake topology. The group $Sp(2n, \mathbb{Q})$ acts on $\overline{\mathfrak{h}}_n$, the involution $\tau : \mathfrak{h}_n \rightarrow \mathfrak{h}_n$ extends to $\overline{\mathfrak{h}}_n$, and $\tau(gx) = \tau(g)\tau(x)$ for any $g \in Sp(2n, \mathbb{Q})$ and every $x \in \overline{\mathfrak{h}}_n$. Denote by $\pi : \overline{\mathfrak{h}}_n \rightarrow \Gamma(4m) \backslash \overline{\mathfrak{h}}_n = V$ the projection to the Baily-Borel compactification of X . The involution τ passes to complex conjugation $\tau : V \rightarrow V$, whose fixed points we denote by $V_{\mathbb{R}}$. Clearly $X_{\mathbb{R}} \subset V_{\mathbb{R}}$. Define $\overline{X}_{\mathbb{R}}$ to be the closure of $X_{\mathbb{R}}$ in $V_{\mathbb{R}}$.

In [Sh], Shimura shows that the $\Gamma(N)$ -automorphic forms on $\mathfrak{h}_n$ are generated by those automorphic forms with rational Fourier coefficients. It follows that:

8.1. Theorem. *There exists a natural rational structure on the Baily-Borel compactification V of X which is compatible with the τ -real structure.*

The Baily-Borel compactification V is stratified by finitely many strata of the form $\pi(F)$ where $F \subset \overline{\mathfrak{h}}_n$ is a rational boundary component. Each such F is isomorphic to some Siegel space $\mathfrak{h}_k$, in which case we say the stratum $\pi(F)$ is a boundary stratum of rank k . Let V^r denote the union of all boundary strata of rank $n - r$. In Proposition 9.2 we will prove that

$$V_{\mathbb{R}}^0 \cup V_{\mathbb{R}}^1 \subset \overline{X}_{\mathbb{R}} \subset V_{\mathbb{R}} \quad (8.1.1)$$

where $V_{\mathbb{R}}^r = V^r \cap V_{\mathbb{R}}$.

As in §3, if F is a rational boundary component of $\mathfrak{h}_n$ which is preserved by the involution τ , and if $\gamma \in \Gamma(4m)$, we define the set of γ -real points of F to be

$$F^\gamma = \{x \in F \mid \tau(x) = \gamma x\}.$$

Then $\pi(F^\gamma) \subset V_{\mathbb{R}}$. A $\Gamma(4m)$ -real boundary pair (F, γ) (of rank q) consists of a rational boundary component F (of rank q) and an element $\gamma \in \Gamma(4m)$ such that $F^\gamma \neq \emptyset$. (Hence $\tilde{F} = \gamma F$.) We say two real boundary pairs (F, γ) and (F_1, γ_1) are equivalent if the resulting locus of real points $\pi(F^\gamma) = \pi(F_1^{\gamma_1})$ coincide. If (F, γ) is a real boundary pair and if $g \in \Gamma(4m)$ then $(gF, \tilde{g}\gamma g^{-1})$ is an equivalent real boundary pair.

8.2. Standard boundary components. Fix an integer q with $1 \leq q < n$. The Siegel upper halfspace

$$\mathfrak{h}_q = \{Z \in M_{q \times q}(\mathbb{C}) \mid {}^t Z = Z \text{ and } \text{Im}(Z) > 0\}$$

is attached to $\mathfrak{h}_n$ as a limit of matrices in $M_{n \times n}(\mathbb{C})$ by

$$Z \mapsto \lim_Y \begin{pmatrix} Z & 0 \\ 0 & iY \end{pmatrix}.$$

See (2.2.2). Here, $Y \in C_{n-q}$ is a positive definite symmetric matrix of order $n - q$, and the limit is taken as $Y \rightarrow \infty C_{n-q}$. This means that for any $Y_0 \in C_{n-q}$, the difference $Y - Y_0$ is eventually contained in C_{n-q} (or, alternatively, that all the eigenvalues of Y converge to ∞). Denote this mapping by $\phi : \mathfrak{h}_q \rightarrow \bar{\mathfrak{h}}_n$. Its image $F_q = \phi(\mathfrak{h}_q) \subset \bar{\mathfrak{h}}_n$ is called the *standard boundary component* of (maximal) rank q . The normalizer in $Sp(2n, \mathbb{Z})$ of F_q is the parabolic subgroup

$$P_q = \left(\begin{array}{cc|cc} A & 0 & B & * \\ * & * & * & * \\ \hline C & 0 & D & * \\ 0 & 0 & 0 & * \end{array} \right)$$

with unipotent radical

$$\mathcal{U}(P_q) = \left(\begin{array}{cc|cc} I_q & 0 & 0 & b \\ a & I_{n-q} & {}^t b & d \\ \hline 0 & 0 & I_q & -{}^t a \\ 0 & 0 & 0 & I_{n-q} \end{array} \right)$$

and Levi factor $L(P_q) = G_h G_\ell$ with

$$G_h = \left(\begin{array}{cc|cc} A & 0 & B & 0 \\ 0 & I_{n-q} & 0 & 0 \\ \hline C & 0 & D & 0 \\ 0 & 0 & 0 & I_{n-q} \end{array} \right) \text{ and } G_\ell = \left(\begin{array}{cc|cc} I_q & 0 & 0 & 0 \\ 0 & T & 0 & 0 \\ \hline 0 & 0 & I_q & 0 \\ 0 & 0 & 0 & {}^t T^{-1} \end{array} \right) \quad (8.2.1)$$

being the “Hermitian” and “linear” factors respectively, where $\begin{pmatrix} A & B \\ C & D \end{pmatrix} \in Sp(2m, \mathbb{R})$ and $T \in GL_{n-q}(\mathbb{R})$. The subgroup $\mathcal{U}(P_q)G_\ell$ is normal in P_q and we denote by

$$\nu : P_q \rightarrow G_h \cong \mathbf{Sp}(2q, \mathbb{R})$$

the projection to the quotient. Then ν commutes with the involution τ .

The boundary component F_q is preserved by τ . The set

$$F_q^\tau = \phi(iC_q) = \{\phi(iY) \mid Y > 0\}$$

of I -real points in F_q is just the set of τ -fixed points in F_q and it may be canonically identified with the cone of positive definite matrices of order q . Denote by iI_q its canonical basepoint.

The boundary component F_q is attached to $\mathfrak{h}_n$ so that this cone $\phi(iC_q)$ is contained in the closure of the cone iC_n .

8.3. Proposition. *Let (F, γ) be a $\Gamma(4m)$ -real boundary pair of rank q . Then there exists $a \in \mathbf{Sp}(2n, \mathbb{Z})$ so that $a(F_q) = F$ and*

$$\tilde{a}^{-1}\gamma a = \begin{pmatrix} A & B \\ 0 & {}^tA^{-1} \end{pmatrix} \in \ker(\nu). \quad (8.3.1)$$

Moreover, we may take $B = 0$, that is, there exists $\gamma' \in \Gamma(4m)$ and $g \in \mathbf{Sp}(2n, \mathbb{Z})$ so that $F^{\gamma'} = F^\gamma$, $g(F_q) = F$, and so that

$$\tilde{g}^{-1}\gamma'g = \begin{pmatrix} A & 0 \\ 0 & {}^tA^{-1} \end{pmatrix} \in \ker(\nu). \quad (8.3.2)$$

8.4. Proof. Choose $b \in \mathbf{Sp}(2n, \mathbb{Z})$ so that $bF_q = F$. Then $w = \tilde{b}^{-1}\gamma b = \tilde{b}^{-1}b \cdot b^{-1}\gamma b \in \Gamma(2)$. Since it also preserves the standard boundary component F_q , we have: $w \in P_q$.

The Hermitian part $\nu(w) \in \mathbf{Sp}(2n, \mathbb{Z})$ acts on the standard boundary component $F_q \cong \mathfrak{h}_q$. By Lemma 4.5, $\nu(\tilde{w})\nu(w) \in \Gamma(4)$ which is torsion-free. By assumption, the set of $\nu(w)$ -real points $F_q^{\nu(w)}$ is nonempty. Choose a point $x \in F_q^{\nu(w)}$ whose stabilizer in $\mathbf{Sp}(2m, \mathbb{Z})$ consists of $\pm I$: by Proposition 3.13 such points exist and are even dense in $F_q^{\nu(w)}$. Then, by Proposition 4.7 there exists $h \in \mathbf{Sp}(2m, \mathbb{Z})$ such that $\tilde{h}h^{-1} = \nu(w)$ and $F_q^{\nu(w)} = h(iC_q)$. Let us identify the element h with its image in $\mathbf{Sp}(2n, \mathbb{Z})$ as in (8.2.1) and set $v = \tilde{h}^{-1}wh$. Then $\nu(v) = 1$. The following diagram may help to sort out these various transformations.

$$\begin{array}{ccccc} F_q & \xrightarrow{\tilde{h}} & F & \xrightarrow{\tilde{b}} & F \\ v \uparrow & & \uparrow w & & \uparrow \gamma \\ F_q & \xrightarrow{h} & F & \xrightarrow{b} & F \end{array}$$

Then the element $a = bh \in \mathbf{Sp}(2n, \mathbb{Z})$ has the desired properties, that is, $a \cdot F_q = F$ and $\nu(v) = 1$ where $v = \tilde{a}^{-1}\gamma a \in \Gamma(2)$.

Now let us prove the “moreover” part of Proposition 8.3. By Lemma 4.5, both $\tilde{\gamma}\gamma^{-1}$ and $\tilde{\gamma}\gamma$ are in $\Gamma(8m)$. Then $\tilde{\gamma}\gamma = \tilde{a}\tilde{v}v\tilde{a}^{-1} \in \Gamma(8m)$, hence $\tilde{v}v \in \Gamma(8m)$. Calculating $\tilde{v}v \equiv I \pmod{8m}$ gives

$$AB \equiv B {}^tA^{-1} \pmod{8m} \text{ and } A^2 \equiv I \pmod{8m}. \quad (8.4.1)$$

Since $v = \tilde{a}^{-1}aa^{-1}\gamma a \in \Gamma(2)$ the matrix A is integral and B is even, so

$$x = \begin{pmatrix} I & -\frac{1}{2}A^{-1}B \\ 0 & I \end{pmatrix}$$

is integral. Since $v \in \ker(\nu)$ we see that $A = \begin{pmatrix} * & 0 \\ * & * \end{pmatrix}$ and $B = \begin{pmatrix} 0 & * \\ * & * \end{pmatrix}$ from which it follows that $x \in \mathcal{U}(P_q) \subset \ker(\nu)$. Set $g = ax$ and set $u' = \tilde{x}^{-1}vx$. Direct computation with the matrices

for x and v gives

$$u' = \begin{pmatrix} A & B' \\ 0 & {}^tA^{-1} \end{pmatrix}$$

where $B' = \frac{1}{2}B - \frac{1}{2}A^{-1}B {}^tA^{-1}$. Using (8.4.1) gives $B' \equiv 0 \pmod{4m}$. The following diagram may help to explain these transformations.

$$\begin{array}{ccccc} F_q & \xrightarrow{\tilde{x}} & F_q & \xrightarrow{\tilde{a}} & F \\ \uparrow u' & & \uparrow v & & \uparrow \gamma \\ F_q & \xrightarrow{x} & F_q & \xrightarrow{a} & F \end{array}$$

Now decompose $u' = uu_2$ where

$$u = \begin{pmatrix} A & 0 \\ 0 & {}^tA^{-1} \end{pmatrix} \text{ and } u_2 = \begin{pmatrix} I & A^{-1}B' \\ 0 & I \end{pmatrix} \in \ker(\nu).$$

Set $\gamma' = \tilde{g}ug^{-1}$. We will verify that g and γ' satisfy the conclusions of Lemma 8.3.

First note that $u_2 \in \Gamma(4m)$ since $B' \equiv 0 \pmod{4m}$. Then $\gamma' \in \Gamma(4m)$ because

$$\gamma = \tilde{a}va^{-1} = \tilde{g}u'g^{-1} = (\tilde{g}ug^{-1})(gu_2g^{-1}) = \gamma'(gu_2g^{-1}) \in \Gamma(4m).$$

We have already verified that $gF_q = F$ and that $u = \tilde{g}^{-1}\gamma'g$ has the desired form. Since u_2 acts trivially on F_q we see that $y \in F^{\gamma'}$ iff

$$\tilde{y} = \gamma'y = \tilde{g}ug^{-1}y = \tilde{g}uu_2g^{-1}y = \gamma y$$

iff $y \in F^\gamma$. Hence $F^{\gamma'} = F^\gamma$. □

8.5. Some nearby boundary components. Fix r with $q \leq r \leq n$ and set $s = n - r$. Define

$$j_r = \left(\begin{array}{cc|cc} I_r & 0 & 0 & 0 \\ 0 & 0 & 0 & I_s \\ \hline 0 & 0 & I_r & 0 \\ 0 & -I_s & 0 & 0 \end{array} \right) \text{ so } j_r^2 = \left(\begin{array}{cc|cc} I_r & 0 & 0 & 0 \\ 0 & -I_s & 0 & 0 \\ \hline 0 & 0 & I_r & 0 \\ 0 & 0 & 0 & -I_s \end{array} \right).$$

Then $\tilde{j}_r = j_r^{-1}$. Although $j_r \notin P_q$, its square j_r^2 preserves F_q and in fact it acts as the identity on F_q . Let $E_{q,r} = j_r(F_q) = j_r^{-1}(F_q)$ and let $Q_{q,r}$ be its normalizing maximal parabolic subgroup: it is j_r -conjugate to P_q . (If $r = n$ then j_r is the identity and $E_{q,r} = F_q$). The involution τ preserves the boundary component $E_{q,r}$ and the set of τ -fixed points

$$E_{q,r}^\tau = \{x \in E_{q,r} \mid \tilde{x} = x\} = j_r \cdot F_q^\tau$$

is contained in the closure of $\mathfrak{h}_n^\tau = iC_n$ as follows. Although j_r does not preserve the cone iC_n , it does preserve the sub-cone $S_{q,r} \subset iC_n$ of elements

$$\begin{pmatrix} iY_1 & 0 & 0 \\ 0 & iY_2 & 0 \\ 0 & 0 & iY_3 \end{pmatrix} \in \mathfrak{h}_n$$

where $Y_1 \in C_q$, $Y_2 \in C_{r-q}$, and $Y_3 \in C_s$. Moreover, F_q^τ is contained in the closure of $S_{q,r}$. Therefore

$$E_{q,r}^\tau = j_r F_q^\tau \subset j_r \overline{S_{q,r}} = \overline{S_{q,r}} \subset \overline{iC_n}. \quad (8.5.1)$$

8.6. Proposition. *Let (F, γ) be a $\Gamma(4m)$ -real boundary pair of rank q . Let $g \in \mathbf{Sp}(2n, \mathbb{Z})$. Suppose that $g(F_q) = F$ and that $u = \tilde{g}^{-1}\gamma g \in \ker(\nu)$. Suppose also that there exists r with $q \leq r \leq n$ so that*

$$\tilde{j}_r^{-1} u j_r = j_r u j_r \in \Gamma(4m).$$

Define $\omega = \tau(gj_r)(gj_r)^{-1}$. Then $\omega \in \Gamma(4m)$ and

$$F^\gamma = F^\omega \subset \overline{\mathfrak{h}_n^\omega}.$$

Consequently the resulting set $\pi(F^\gamma)$ of real points is contained in the closure $\overline{X}_{\mathbb{R}}$.

8.7. Proof. Calculate

$$\omega = \tau(gj_r)(gj_r)^{-1} = \gamma((gj_r)(j_r u j_r)^{-1}(gj_r)^{-1}) \in \Gamma(4m)$$

(using the fact that $\tilde{j}_r = j_r^{-1}$) which proves the first statement. Since $u \in P_q$ acts trivially on F_q , the same is true of $j_r^2 u$. Hence, $x \in F^\omega$ if and only if

$$\tilde{x} = \omega x = \tau(gj_r)j_r^{-1}(g^{-1}x) = \tau(gj_r)j_r^{-1}j_r^2 u(g^{-1}x) = \tau(gj_r)\tau(gj_r)^{-1}\gamma x = \gamma x$$

which holds if and only if $x \in F^\gamma$. The following diagram may help in placing these elements,

$$\begin{array}{ccccc} E_{q,r} & \xrightarrow{\tilde{j}_r} & F_q & \xrightarrow{\tilde{g}} & F \\ \uparrow & & \uparrow u & & \uparrow \gamma \\ E_{q,r} & \xrightarrow{j_r} & F_q & \xrightarrow{g} & F \end{array}$$

We claim that $F^\omega = gj_r(E_{q,r}^\tau)$. In fact, $x \in E_{q,r}^\tau$ if and only if $\tilde{x} = x$ which holds iff

$$\tau(gj_r)\tilde{x} = \tau(gj_r)x = \tau(gj_r)(gj_r)^{-1}(gj_r)x = \omega(gj_r)x$$

which holds iff $(gj_r)x \in F^\omega$. Similarly, $gj_r(\mathfrak{h}_n^\tau) = \mathfrak{h}_n^\omega$. Hence

$$F^\omega = gj_r(E_{q,r}^\tau) \subset gj_r \overline{\mathfrak{h}_n^\tau} = \overline{\mathfrak{h}_n^\omega}$$

using (8.5.1). □

9. CORANK ONE STRATA

9.1. In this section we specialize to the case $q = n-1$, that is, we consider only the boundary strata of maximal rank. As in §8, F_{n-1} denotes the standard boundary component of corank 1, P_{n-1} denotes its normalizing parabolic subgroup and G_h and G_ℓ refer to the Hermitian and linear factors (8.2.1) of the Levi quotient $L(P_{n-1})$.

9.2. Theorem. *Let F be a proper rational boundary component of $\mathfrak{h}_n$ with (maximal) rank $n-1$, let $\gamma \in \Gamma(4m)$, and suppose that $F^\gamma \neq \phi$. Then F^γ is contained in the closure of the set $\mathfrak{h}_n^{\Gamma(4m)}$ of $\Gamma(4m)$ -real points of $\mathfrak{h}_n$.*

9.3. **Proof.** By Proposition 8.3, there exists $\gamma' \in \Gamma(4m)$ and $g \in \mathbf{Sp}(2n, \mathbb{Z})$ so that $gF_{n-1} = F$, $F^\gamma = F^{\gamma'}$ and so that

$$u = \tilde{g}^{-1}\gamma'g = \begin{pmatrix} A & 0 \\ 0 & {}_tA^{-1} \end{pmatrix} \in \ker(\nu).$$

This implies that $A = \begin{pmatrix} I_{n-1} & 0 \\ a & \pm 1 \end{pmatrix}$. Note that $\tilde{u}u = g^{-1}\tilde{\gamma}\gamma g \in \Gamma(8m)$ by Lemma 4.5. Hence $a \equiv 0 \pmod{4m}$.

If the plus sign occurs then this says that $u \in \Gamma(4m)$. Let $\omega = \tilde{g}g^{-1}$. Then Proposition 8.6 (with $q = n-1$ and $r = n$) implies that $F^{\gamma'} \subset \overline{\mathfrak{h}_n^\omega}$.

If the minus sign occurs then

$$j_{n-1}uj_{n-1} = \left(\begin{array}{cc|cc} I & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ \hline 0 & -{}_t a & I & 0 \\ -a & 0 & 0 & 1 \end{array} \right) \in \Gamma(4m)$$

so we may apply Proposition 8.6 (with $q = r = n-1$) to conclude that $F^{\gamma'} \subset \overline{\mathfrak{h}_n^\omega}$ where $\omega = \tau(gj_{n-1})(gj_{n-1})^{-1}$. $\square$

10. THE PRINCIPAL CONGRUENCE GROUP $\Gamma(2^k)$

10.1. Throughout this section we let $\Gamma = \Gamma(2^k) \subset \mathbf{Sp}(2n, \mathbb{Z})$ be the principal congruence subgroup of level 2^k with $k \geq 2$. As in §8, let $X = X_{\mathbb{C}} = \Gamma \backslash \mathfrak{h}_n$ and let $V = V_{\mathbb{C}} = \Gamma \backslash \overline{\mathfrak{h}_n}$ be its Baily-Borel compactification with projection $\pi : \overline{\mathfrak{h}_n} \rightarrow V$. Let $\overline{X}_{\mathbb{R}}$ be the closure of $X_{\mathbb{R}}$ in V . In this section we will prove that $\overline{X}_{\mathbb{R}} = V_{\mathbb{R}}$. The proof of the following lemma will appear in §10.7.

10.2. Lemma. *Let $A \in \mathbf{GL}(n, \mathbb{Z})$. Suppose that*

$$A \equiv I \pmod{2} \text{ and } A^2 \equiv I \pmod{2^{k+1}}.$$

Then there exists $p \in \mathbf{GL}(n, \mathbb{Z})$ so that

$$p^{-1}Ap \equiv \begin{pmatrix} \pm 1 & & & \\ & \pm 1 & \dots & \\ & & \dots & \\ & & & \pm 1 \end{pmatrix} \pmod{2^k}. \tag{10.2.1}$$

Moreover if the matrix of A with respect to the standard basis of $\mathbb{Z}^n$ is $\begin{pmatrix} I_q & 0 \\ * & * \end{pmatrix}$ then it is possible to choose p to be of the form $p = \begin{pmatrix} I_q & 0 \\ * & * \end{pmatrix}$.

10.3. Lemma. *Let (F, γ) be a real boundary pair of rank q . Then there exists $r \geq q$, there exists $\gamma' \in \Gamma(2^k)$ and there exists $g \in \mathbf{Sp}(2n, \mathbb{Z})$ such that $F^\gamma = F^{\gamma'}$, $gF_q = F$, $\tilde{g}^{-1}\gamma'g \in \ker(\nu)$, and so that*

$$\tilde{g}^{-1}\gamma'g \equiv \left(\begin{array}{cc|cc} I_r & 0 & 0 & 0 \\ 0 & -I_s & 0 & 0 \\ \hline 0 & 0 & I_r & 0 \\ 0 & 0 & 0 & -I_s \end{array} \right) \pmod{2^k} \quad (10.3.1)$$

where $s = n - r$.

10.4. Proof. By Proposition 8.3 there exists $\gamma' \in \Gamma(2^k)$ and there exists $a \in \mathbf{Sp}(2n, \mathbb{Z})$ so that $F^\gamma = F^{\gamma'}$, $aF_q = F$, and

$$u = \tilde{a}^{-1}\gamma'a = \begin{pmatrix} A & 0 \\ 0 & {}_tA^{-1} \end{pmatrix} \in \ker(\nu).$$

Then $\tilde{\gamma}'\gamma' = au^2a^{-1} \in \Gamma(2^{k+1})$ by Lemma 4.5, hence $A^2 \equiv I \pmod{2^{k+1}}$. Moreover, $A \in \Gamma(2)$ and $A = \begin{pmatrix} I_q & 0 \\ * & * \end{pmatrix}$. Let $p \in \mathbf{GL}(n, \mathbb{Z})$ be the change of basis provided by Lemma 10.2. Then $p^{-1}Ap = \begin{pmatrix} I_q & 0 \\ * & * \end{pmatrix}$ and (after re-ordering the coordinates if necessary), $p^{-1}Ap \equiv \begin{pmatrix} I_r & 0 \\ 0 & -I_s \end{pmatrix} \pmod{2^k}$ for some $r \geq q$. Set $h = \begin{pmatrix} p & 0 \\ 0 & {}_tp^{-1} \end{pmatrix} \in \ker(\nu)$. Set $g = ah$.

$$\begin{array}{ccccc} F_q & \xrightarrow{\tilde{h}} & F_q & \xrightarrow{\tilde{a}} & F \\ \uparrow & & \uparrow u & & \uparrow \gamma' \\ F_q & \xrightarrow{h} & F_q & \xrightarrow{a} & F \end{array}$$

Then $\tilde{h}^{-1}uh = \tilde{g}^{-1}\gamma'g \in \ker(\nu)$ and $\tilde{g}^{-1}\gamma'g$ has the desired form (10.3.1). $\square$

10.5. Theorem. *Let (F, γ) be a real boundary pair. Then there exists $\gamma_1 \in \Gamma(2^k)$ so that the set $F^\gamma = F^{\gamma_1}$ of γ -real points is contained in the closure $\overline{\mathfrak{h}_n^{\gamma_1}}$.*

10.6. Proof. Set $q = \text{rank}(F)$. By Lemma 10.3, there exists $\gamma_1 \in \Gamma(2^k)$ and there exists $g \in \mathbf{Sp}(2n, \mathbb{Z})$ so that $F^\gamma = F^{\gamma_1}$, so that $g(F_{n-1}) = F$ and so that $u = \tilde{g}^{-1}\gamma_1g$ lies in $\ker(\nu)$ and has the form (10.3.1), for some $r \geq q$. Therefore $j_ru_jr \equiv I \pmod{2^k}$ so Proposition 8.6 may be applied. $\square$

10.7. Proof of Lemma 10.2. The lemma is equivalent to the following statement. Suppose M is a free $\mathbb{Z}$ -module of rank n . Let $\alpha : M \rightarrow M$ be an automorphism such that $(\alpha - I)M \subset 2M$ and $(\alpha^2 - I)M \subset 2^{k+1}M$, that is, $\alpha \equiv I \pmod{2}$ and $\alpha^2 \equiv I \pmod{2^{k+1}}$. Then there exists a basis $x_1, x_2, \dots, x_n$ of M so that $\alpha(x_i) = \pm x_i \in 2^k M$ for $i = 1, 2, \dots, n$.

This statement will be proven by induction on the rank of M . The case of rank 1 is obvious, so suppose that M has rank n . We will show that there exists a basis $\{x_1, x_2, \dots, x_n\}$ of M so that $\alpha(x_1) \equiv \pm x_1 \pmod{2^k}$. If $\ker(\alpha - I)$ is not trivial or if $\ker(\alpha + I)$ is not trivial then any primitive x within this kernel may be extended to a basis. Therefore we may assume that both $(\alpha - I)M$ and $(\alpha + I)M$ have maximal rank.

By elementary divisor theory there exists a basis $x_1, x_2, \dots, x_n$ of M and integers $d_1, d_2, \dots, d_n$ so that $d_1 | d_2 | \dots | d_n$ and so that $d_1 x_1, d_2 x_2, \dots, d_n x_n$ is a basis of $(\alpha - I)M$. Similarly there exists another basis $y_1, y_2, \dots, y_n$ of M and integers $e_1 | e_2 | \dots | e_n$ so that $e_1 y_1, e_2 y_2, \dots, e_n y_n$ is a basis for the submodule $(\alpha + I)M \subset M$.

We claim that either $\alpha(x_1) \equiv -x_1 \pmod{2^k}$ or $\alpha(y_1) \equiv y_1 \pmod{2^k}$. First note that $(\alpha + I)d_1 x_1 \equiv 0 \pmod{2^{k+1}}$ since $d_1 x_1 \in (\alpha - I)M$. If d_1 is odd this implies $\alpha x_1 + x_1 \equiv 0 \pmod{2^{k+1}}$. If $d_1/2$ is odd, it implies that $\alpha x_1 + x_1 \equiv 0 \pmod{2^k}$. Similarly, if e_1 is odd or if $e_1/2$ is odd then $\alpha y_1 - y_1 \equiv 0 \pmod{2^k}$. However, $(\alpha - I)M + (\alpha + I)M = 2M$ so the highest power of 2 which divides $\gcd(d_1, e_1)$ is 2^1 . Therefore one of these four cases must occur, which proves the claim.

By switching the x 's with the y 's if necessary, we arrive at a basis $x_1, x_2, \dots, x_n$ of M such that $\alpha(x_1) \equiv \pm x_1 \pmod{2^k}$. Write $M = M_1 \oplus M_2$ where $M_1 = \mathbb{Z}x_1$ and $M_2 = \sum_{i \geq 2} \mathbb{Z}x_i$. With respect to this decomposition, α has the matrix $\begin{pmatrix} \alpha_{11} & \alpha_{12} \\ \alpha_{21} & \alpha_{22} \end{pmatrix}$ where $\alpha_{11} \equiv \pm 1 \pmod{2^k}$, and where $\alpha_{22} : M_2 \rightarrow M_2$. We claim that $\alpha_{22} \equiv I \pmod{2}$ and $\alpha_{22}^2 \equiv I \pmod{2^{k+1}}$. Since $\alpha \equiv I \pmod{2}$ we have

$$\alpha \begin{pmatrix} 0 \\ m_2 \end{pmatrix} = \begin{pmatrix} \alpha_{12} m_2 \\ \alpha_{22} m_2 \end{pmatrix} \equiv \begin{pmatrix} 0 \\ m_2 \end{pmatrix} \pmod{2}$$

hence $\alpha_{12} \equiv 0 \pmod{2}$ and $\alpha_{22} \equiv I \pmod{2}$. Also,

$$\alpha \begin{pmatrix} x_1 \\ 0 \end{pmatrix} = \begin{pmatrix} \alpha_{11} x_1 \\ \alpha_{21} x_1 \end{pmatrix} \equiv \begin{pmatrix} \pm x_1 \\ 0 \end{pmatrix} \pmod{2^k}$$

hence $\alpha_{21} \equiv 0 \pmod{2^k}$. Similarly,

$$\alpha^2 \begin{pmatrix} 0 \\ m_2 \end{pmatrix} = \begin{pmatrix} * \\ \alpha_{21} \alpha_{12} m_2 + \alpha_{22}^2 m_2 \end{pmatrix} \equiv \begin{pmatrix} 0 \\ m_2 \end{pmatrix} \pmod{2^{k+1}}.$$

But $\alpha_{12} \equiv 0 \pmod{2}$ and $\alpha_{21} \equiv 0 \pmod{2^k}$ so the first term in this sum is congruent to 0 $\pmod{2^{k+1}}$, hence $\alpha_{22}^2 m_2 \equiv m_2 \pmod{2^{k+1}}$ as claimed.

Therefore we may apply induction to the pair (M_2, α_{22}) to obtain a basis, which we again denote by $x_2, x_3, \dots, x_n$ such that $\alpha_{22}(x_j) \equiv \pm x_j \pmod{2^k}$ for $j \geq 2$. Hence, for $j \geq 2$, there are integers a_j and a sign $\epsilon_j = \pm 1$ so that $\alpha(x_j) \equiv a_j x_1 + \epsilon_j x_j \pmod{2^k}$. Each a_j is even since $\alpha \equiv I \pmod{2}$. Define a new basis

$$x'_j = \begin{cases} x_j & \text{if } \epsilon_j = \epsilon_1 \\ \epsilon_j x_j + \frac{1}{2} a_j x_1 & \text{if } \epsilon_j = -\epsilon_1. \end{cases}$$

We claim this basis has the desired property: $\alpha(x'_j) = \pm x'_j \pmod{2^k}$. First suppose $\epsilon_j = +\epsilon_1$. Then $\alpha^2(x_j) = 2a_j\epsilon_j x_1 + x_j$. Since $\alpha^2 \equiv I \pmod{2^{k+1}}$ we see that $a_k \equiv 0 \pmod{2^k}$. Hence $\alpha(x'_j) \equiv x_j \pmod{2^k}$ as desired. If $\epsilon_j\epsilon_1 = -1$ then

$$\begin{aligned}\alpha(x'_j) &= \epsilon_j\alpha(x_j) + \frac{1}{2}a_j\alpha(x_1) \\ &\equiv \epsilon_j(a_j x_1 + \epsilon_j x_j) + \frac{1}{2}a_j\epsilon_1 x_1 \pmod{2^k} \\ &\equiv x_j + \frac{1}{2}\epsilon_j a_j x_1 \pmod{2^k} \\ &= \epsilon_j(\epsilon_j x_j + \frac{1}{2}a_j x_1) = \epsilon_j x'_j.\end{aligned}$$

This completes the construction of the desired basis.

To prove the “moreover” part of the lemma, let $M' = \sum_{i=1}^q \mathbb{Z}e_i$ be the submodule of M generated by the first q standard basis vectors. Apply the lemma to the quotient module $\alpha : M/M' \rightarrow M/M'$. Choose any lift $x_{q+1}, \dots, x_n$ of the resulting basis of M/M' to M and define $x_1 = e_1, x_2 = e_2, \dots, x_q = e_q$. With respect to this basis,

$$\alpha(x_j) = \epsilon_j x_j + \sum_{i=1}^q a_{ij} x_i \text{ for } i > q$$

where $\epsilon_i = \pm 1$ and a_{ij} is even. Set

$$x'_i = \begin{cases} x_i & \text{if } \epsilon_i = 1 \\ -x_i + \sum_{j=1}^q \frac{1}{2}a_{ij}x_j & \text{if } \epsilon_i = -1 \end{cases}.$$

Then $\alpha(x'_i) \equiv \epsilon_i x'_i \pmod{2^k}$ and the change of basis matrix has the desired form. $\square$

REFERENCES

- [C] H. Comessatti, Sulle varietà abeliane reali I, II. Ann. Mat. Pura. Appl. **2** (1924) pp. 67-106 and **4** (1926) pp. 27-72.
- [La] H. Lange and C. Birkenhake, **Complex Abelian Varieties**, Grundlehren Math. Wiss. **302**, Springer Verlag, Berlin, 1992.
- [Si] R. Silhol, Compactifications of moduli spaces in real algebraic geometry. Inv. Math. **107** (1992), pp. 151-202.
- [Si2] R. Silhol, Real abelian varieties and the theory of Comessatti. Math. Z. **181** (1982), pp. 345-362.
- [Sh] G. Shimura, On the Fourier coefficients of modular forms of several variables. Nachr. Akad. Wiss. Göttingen Math-Phys. **17** (1975), 261-268.